

// ()
 // , ,
 //

:

*

:

()
 () :
 ()

Pure Tone Audiometry

Relative Risk

Chi-square

$\pm / / \pm /$:
 RR (P< /) % % (P< /)
 / RR (P< /) % % /
 :

:

()

()

()

()

()

Historical Cohort Study

.()

()

.()

(Sound Level Meter)

() ()

()

()

.()

()

(())

()

%

(Drop out) (Pure Tone Audiometry) PTA

(t = / df = P < /)

(P < /)

(% ,)

t-test

Chi Square
(Relative Risk)

(P < /)

(CI)

%

()

(Pilot)

%

%

P2 P1

Chi Square

%

(% %)

(P < /)

%

(P< /) % (Relative Risk)
(Relative Risk)

/ (% CI = / - /)
(% CI = / - /) %

()	()	()	()	
± SD	± SD	± SD	± SD	
/ ± /	/ ± /	/ ± /	±	
/ ± /	/ ± /	/ ± /	/ ±	
P< /	P< /	P< /	P< /	

()	()	*()	
± SD	± SD	± SD	
/ ± /	/ ± /	/ ± /	
/ ± /	/ ± /	/ ± /	

(t=2.1, df= 118, p<0.05) -*

()				
()	()	()	()	
()	()	()	()	

()

)

(

Lusk

()

()

()

.()

.

.

()

Santana

()

.()

.

.

.(-)

Palmer

/

.()

Palmer

.()

.()

/

- : ()

()

REFERENCES

1. Clark W. Hearing: the effect of noise. *Otolaryngol Head Neck Surg* 1992; 106(6): 669-76.
2. Landen D, Wilkins S, Staphenson M, McWilliams L. Noise exposure and hearing loss among sand and gravel miners. *Occup Environ Hyg* 2004; 1(8): 532-41.
3. Osguthorpe JD, Klein AJ. Occupational hearing conservation. *Otolaryngol Clin North Am.* 1991; 24(2): 403-14.

-
5. Guidotti TL, Novak RE. Hearing conservation and occupational exposure to noise. *Am Fam Physician* 1983; 28(4): 181-6.
 6. Baker DB, Landrigan PJ. Occupationally related disorders. *Med Clin North Am* 1990; 74(2): 441-60.
 7. Williams W, Purdy S, Murray N, LePage E, Challinor K. Hearing loss and perceptions of noise in the workplace among rural Australians. *Aust J Rural Health* 2004; 12(3):115-9.
 9. Lusk SL, Hagerty BM, Gillespie B, Caruso CC. Chronic effects of workplace noise on blood pressure and heart rate. *Arch Environ Health* 2002; 57(4): 273-81.
 10. Santana VS, Barberino JL. Occupational noise exposure and hypertension. *Rev Saude Publica*. 1995, 29(6): 478-87.
 11. Reilly MJ, Rosenman KD, Kalinowski DJ. Occupational noise-induced hearing loss surveillance in Michigan. *J Occup Environ Med* 1998; 40(8): 667-74.
 12. Palmer KT, Griffin MJ, Syddall HE, Davis A, Pannett B, Coggon D. Occupational exposure to noise and the attributable burden of hearing difficulties in Great Britain. *Occup Environ Med* 2002; 59: 634-9.
 13. Palmer KT, Griffin MJ, Syddall HE, Coggon D. Cigarette smoking, occupational exposure to noise, and self reported hearing difficulties. *Occup Environ Med* 2004; 61: 340-4.
 14. Irwin J. What are the causes, prevention and treatment of hearing loss in the ageing worker? *Occup Med* 2000; 50(7): 492-5.
 15. Cruickshanks KJ, Wiley TL, Tweed TS, Klein BE, Klein R, Mares-Perlman JA, et al. Prevalance of hearing loss in older adults in Beaver Dam, Wisconsin. The epidemiology of hearing loss study. *Am J Epidemiol* 1998; 148(9): 879-86.
 16. Uimonen S, Maki-Torkko E, Sorri M. Hearing and occupation. *Int J Circumpolar Health* 1998; 57(2-3): 156-61.
 17. Saiyed HN, Tiwari RR. Occupational health research in India. *Ind Health* 2004; 42(2): 141-8.