

//
//

()

*

(Temporal Myoplasty)

/ ±
/ ±
% (static)
(dynamic) % % % %
% % %

static

()

()

()

gold standard

()

()

()

()

labbe

caen

(Release)

()

(Coronoid process)

()

(alopecia)

()

(dissection)

- mm



()

()

cm

(temporal crest)




(fossa temporal)

(coronal)

(infra temporal crest)

()

cm



(Nasolabial crease)

(masseter)

(commissure)

/

% %
 % % %
 (dynamic)

% % %

/ ±

)

(

/ ± /

%

/ ± /

%

(static)

***	**	***	***
**	**	***	**
*	**	*	**
***	**	***	***
***	**	***	***
**	**	**	**

*** ** *

() (static) .()
()
()





:



:



:



:

(.)

() gold standard

(bulge)

()

suspension

()

" "

()

()

Masseter

()

()

()

VII - V

(interoral approach)

(commissure)

(occlusal plan)

(myoplasty masseter)

()

Gillis

REFERENCES

1. Zuker RM, Manktelow RT, Hussain G. In: Mathes SJ. Plastic surgery. 2nd ed. Vol 3. Philadelphia: Saunders. 2006 P:883-916.
2. Baker DC. Reconstruction of the Paralysed face. In: Grab and Smith plastic surgery. 5th ed. Philadelphia: Lippincott Raven. 1997; P:545-57.
3. Baker DC. Facial paralysis. In: Mc Carthy Plastic surgery. 1st ed. Philadelphia: W.B Saunders. 1990; P:2255-306.
4. Ueda K, Harii K, Asato H, Yamada A. Neurovascular free muscle transfer combined with cross-face nerve grafting for the treatment of facial paralysis in children. *Plast Reconstr Surg* 1998;101(7):1765-73.
5. May M, Drucker C. Temporalis muscle for facial reanimation. A 13-year experience with 224 procedures. *Arch Otolaryngol Head Neck Surg.* 1993;119(4):378-82; discussion 383-4.
6. Hu ZQ, Ogawa R, Aoki R, Gao JH, Hyakusoku H. Temporalis muscle-galea pedicled flap for reconstruction of longstanding facial paralysis. *J Nippon Med Sch.* 2005;72(2):105-12.
7. Labbé D, Huault M. Lengthening temporalis myoplasty and lip reanimation. *Plast Reconstr Surg.* 2000;105(4):1289-97; discussion 1298.
8. Ueda K, Akiyoshi K, Suzuki Y, Ohkouchi M, Hirose T, Asai E, et al. Combination of hypoglossal-facial nerve jump graft by end-to-side neuroorrhaphy and cross-face nerve graft for the treatment of facial paralysis. *J Reconstr Microsurg.* 2007;23(4):181-7.
9. Alam D. Rehabilitation of long-standing facial nerve paralysis with percutaneous suture-based slings. *Arch Facial Plast Surg.* 2007;9(3):205-9.
10. Garanhani MR, Cardoso JR, Capelli Ade M, Ribeiro MC. Physical therapy in peripheral facial paralysis: retrospective study. *Rev Bras Otorrinolaringol (Engl Ed).* 2007;73(1):106-9.