

//
//

()

, ,

*

, :

"

"

-

- () - :

,

,

Hannalla

:

($P < /$)

,

,

:

,

,

:

:

mohsen_rouzrokh@yahoo.com :

,

,

,

,

,

,

() .()

.()

()

N₂O

.(-)

ACTH

/

.(-)

cc)

(.()

()

Diplus)
Radium

.(

/

—

* TLC: Tender Loving Care

$$(P < \quad)$$
$$(P < \quad)$$

()

()

•

($P < /$)

$$(P = \quad / \quad)$$

(P= /)

$$(P < 1)$$

(Hannalla)

.(P= /)

/ ± /

/ ± /

(P= /)

.(-)

()



*			Mean ± SD	Mean ± SD	Mean ± SD	
/	/	/	/ ± /	/ ± /	/ ± /	μU/ml
			/ ± /	/ ± /	/ ± /	
/	/	/	/ ± /	/ ± /	/ ± /	mg/dL
			/ ± /	/ ± /	/ ± /	
/	/	/	/ ± /	/ ± /	/ ± /	μg/dl
			/ ± /	/ ± /	/ ± /	
/	/	/	/ ± /	/ ± /	/ ± /	mIU/L
			/ ± /	/ ± /	/ ± /	

* Paired t-test (P<0.05)

()



*			Mean ± SD	Mean ± SD	Mean ± SD	
/	/	/	/ ± /	/ ± /	/ ± /	μU/ml
/	< /	< /	/ ± /	/ ± /	/ ± /	
/	< /	< /	/ ± /	/ ± /	/ ± /	mg/dL
/	< /	< /	/ ± /	/ ± /	/ ± /	
/	/	/	/ ± /	/ ± /	/ ± /	μg/dL
/	< /	/	/ ± /	/ ± /	/ ± /	
< /	/	< /	/ ± /	/ ± /	/ ± /	mIU/L
< /	/	< /	/ ± /	/ ± /	/ ± /	

* t-test (P<0.05)



(- ng/ml :)

(> μg/IU/ml :)



(/ - ng/ml :)

(- mg/ml :)

()

()

ACTH

Castejan-Casadoy

ACTH

.()

.()

Anand .()

()

George Sakellaris

()

-

.()

.(-)

()

.()

REFERENCES

1. Denman WT, Ballantyne J. The Massachusetts general Hospital Hand book of pain management. 2nd ed. 2006; part: 22.
2. MC Donald AJ. Cooper Mg: patient-controlled analgesia: An appropriate method of pain control in children. *Paediatr Drugs* 2001; 3: 273-84.
3. Conroy JM, Othersen HB Jr, Dorman BH, Gottesman JD, Wallace CT, Brahen NH. A comparison of wound instillation and caudal block for analgesia following pediatric inguinal hernioaphy. *J Pediatr Surg* 1993; 28: 565-667.
4. Anand KJS, Agnsley-Green A. Measuring the severity of surgical stress in newborn infants. *J Pediatr Surg* 1988; 23: 297-305.
5. Anand KJ, Brown MJ, Causon RC, Christofides ND, Bloom SR, Aynsley-Green A. Can the human neonate mount an endocrine and metabolic response to surgery? *J Pediatr Surg* 1985; 20:41-48
6. Anand KJS Hickey PR. Halothane-morphine compared with high-dose sufentanil for anesthesia and postoperative analgesia in neonatal cardiac surgery. *N Engl J Med* 1992; 326: 1-9.
7. Håkanson E, Rutberg H, Jorfeldt L, Wiklund L. Endocrine and metabolic responses after standardized moderate surgical trauma: Influence of age and sex. *Clin Physiol* 1984; 4: 461-73.
8. Ishida H, Murata N, Yamada H, Nomura T, Shimomura K, Fujioka M, et al. Effect of CO2 pneumoperitoneum on growth of liver micrometastases in a rabbit model. *World J Surg* 2000; 24: 1004-8.
9. Rutberg H, Hakanson E, Anderberg B. Effects of the extradural administration of morphine, or Bupivacaine. *Br J Anasth* 1984; 86: 611-8.
10. Thorell A, Efendic S, Gutniak M, Häggmark T, Ljungqvist O. Insulin resistance after abdominal support in pediatrics. *Nutrition* 1998; 14: 130-42.
11. Casey WF, Rice LJ, Hannallah RS, Broadman L, Norden JM, Guzzetta P. A comparison between bupivacaine instillation versus ilioinguinal/iliohypogastric nerve block for postoperative analgesia following inguinal hernioraphy in children. *Anesthesiology* 1990; 72: 637-9.
12. Ward Platt MP, Tarbit MJ, Aynsley-Green A. The effects of anesthesia and surgery on metabolic homeostasis in infancy and childhood. *J Pediatr Surg* 1990; 25: 472-8.
13. Reid MF, Harris R Phillips RD. Day-case herniotomy in children. A comparison of ilioinguinal nerve block and wound infiltration for postoperative analgesia. *Anaesthesia* 1987; 42: 658-61.

14. Anand KJS. The stress response to surgical trauma: Form physiological basis to therapeutic implications. *Prog Food Nutr Sci* 1986; 10: 67-132.
15. Solak M, Ulusiy H, Sarihan H. Effects of caudal block on cortisol and prolactin responses to postoperative pain in children. *Eur J Pediatr Surg* 2000; 10: 219-23.
16. Okur H, Hucukaydin M, Muhtaroglu S. Effects of bupivacaine on beta-endorphin and cortisol release and postoperative pain following inguinal herniorraphy in children. *Peadiatr Surg Int* 1995; 11: 41-4.
17. Castejón-Casado J, Moreno-Prieto M, Valladares-Mendías JC, Alaminos-Mingorance M, López-Candel E, Ramírez-Navarro A. Hormonal response to sargical stress in school children. *Eur J Pediatr Surg* 2001; 11: 44-7.
18. Reiz S, Haggmark S, Johansson G, Nath S. Cardiotoxicity of ropivacaine: A new amide local anesthesia agent. *Acta Anaesthesiol Scand* 1989; 33(2): 93-8.
19. Wolf AR, Eyres RL, Laussen PC, Edwards J, Stanley IJ, Rowe P, et al: Effect of extradural analgesia on stress response to abdominal surgery in infants. *Br J Anaesth* 1993; 70: 654-60.
20. akellaris G, Petrakis I, Makatounaki K, Arbiros I, Karkavitsas N, Charissis G. Effects of Ropivacaine Infiltration on cortisol and prolactin responses to postoperative pain after inguinal herniorraphy in children. *J Pediatr Surg* 2004; 39(9): 1400-3.
21. Luz G, Innerhofer P, Haussler B, Oswald E, Salner E, Sparr H. Comparison of ropivacaine 0.1% and 0.2% with bupivacaine 0.2% for single-shot caudal anaesthesia in children. *Paediatr Anaesth* 2000, 10:449-504.
22. Taylor MS. Managing of postoperative pain. *Hosp Med* 2001; 62: 560-3.