

## ORIGINAL RESEARCH

### The effectiveness of emotion regulation therapy (ERT) on improving depression, anxiety and stress in patients with myocardial infarction

Bahareh Javadi<sup>1</sup>, Mohamad Hasan Davazdahemamy<sup>\*2</sup>, Mohamad Ismail Gheidari<sup>3</sup>, Mohamad Noori<sup>4</sup>

1. Master of Clinical Psychology, Shahid Beheshti University of Medical Sciences
2. Department of Clinical Psychology, Shahid Beheshti University of Medical Sciences
3. Department of Cardiology, Shahid Beheshti University of Medical Sciences
4. Department of Clinical Psychology, Shahid Beheshti University of Medical Sciences

\*Corresponding Author:

Address: Department of Clinical Psychology, Shahid Beheshti University of Medical Sciences. Tehran, Iran. Tel: +98-21-23031548. Email: mo.emamy@uswr.ac.ir

Date Received: September, 2017    Date Accepted: December, 2017    Online Publication: December 28, 2017

#### Abstract

**Background:** Among the cardiovascular diseases, myocardial infarction is one the main cause of mortality around the world. People with myocardial infarction are significantly more likely to suffer from psychological problems, such as depression, anxiety and stress. Therefore, psychological interventions can help their routine treatment. The purpose of this study was to determine the effectiveness of emotional regulation therapy on depression, anxiety and stress in patients with myocardial infarction.

**Methods:** In this single-case design, 5 patients with myocardial infarction referred to the Taleghani Hospital were selected using convenience sampling method. They received eight sessions of emotional regulation psychotherapy. These patients were evaluated by anxiety, stress, and depression scale (DASS-21) before starting treatment and after each session.

**Results:** The results showed that emotional regulation therapy significantly reduced depression, anxiety and stress in patients with myocardial infarction.

**Discussion:** According to the findings of this study indicating the effectiveness of emotional regulation therapy on depression, anxiety and stress in patients with myocardial infarction, this treatment can be used as part of a comprehensive treatment process for these patients.

**Keywords:** Myocardial infarction, Depression, Anxiety, Stress, Emotional regulation therapy

## Introduction

Myocardial infarction is one of the most common diseases in industrialized and developing countries. In the United States, approximately 525,000 patients each year experience new acute myocardial infarction (AMI), and 190,000 have recurrent AMI. Despite the major advances in primary and secondary prevention of cardiovascular disease, over 1.5 million people experience myocardial infarction every year. By the year 2020, cardiovascular disease is predicted to be the main cause of death and the first cause of disability around the world (1). In the Iranian population, cardiovascular disease is one of the major causes of mortality and disability and is currently the first cause of death in people over the age of 35 in Iran (2).

The heart, as one of the most functional organs, plays a critical role in the health and function of other organs (3); it is obvious that the dysfunction of this organ puts our lives in danger. Several studies have shown that there is a significant relationship between cardiovascular disease and negative emotions such as depression, anxiety and stress. Cardiovascular complications not only include physical symptoms, but also it is significantly associated with psychological disturbances such as anxiety and depression that may lead to changes in the level of performance, occupational status, relationship status and quality of life of individuals (4-6). The meta-analysis that was conducted in 2011 on 29 studies showed that depression increased the risk of mortality associated with cardiovascular disease by 2.7 times in 24 months after myocardial infarction (7). Other studies also have indicated that 45 to 60 percent of patients develop depressive symptoms after a myocardial infarction (8-10). Frasure-Smith (2006) showed that the cost of treatment for depressed patients with cardiovascular disease is 41% higher than non-depressed patients and the cost of treating heart disease will be reduced if the depression is treated (11). Further studies have shown that most patients in the cardiac units have severe anxiety in the first 48 hours, and they consider the main cause of anxiety as fear of dying and fear of recurrent stroke (12, 13). High levels of anxiety and depression will delay the recovery process. Anxiety can increase the Q-T interval (which reflects the tendency to ventricular

irregularities), heart rate, blood pressure and ventricular contractions. Some of the anxiety complications in patients with recurrent myocardial infarction are ventricular tachycardia and ventricular fibrillation. Approximately 70% to 80% of patients with myocardial infarction experience anxiety and this can prevent the adjustment of psychosocial problems and physical recovery after stroke (12, 14). In the case of stress studies, research has shown that the likelihood of a stroke (infraction) after a stressful day can be 9 times higher, and stress can also lead to negative emotions, depression, anxiety, anger, and hostility (15). In general, studies have shown that patients with myocardial infarction experience anxiety long after recovery (remission), and high anxiety may increase the risk of developing myocardial infarction (9, 16); therefore, one of the important issues that patients with myocardial infarction is facing, is their psychological and emotional problems that should be addressed in the treatment process (17). The aim of this study was to investigate the effectiveness of emotional regulation therapy in patients with myocardial infarction. Emotional regulation therapy is an emotion-based therapeutic approach and the purpose of treatment is to improve the patient's psychological function. An important issue in emotional regulation therapy is the patients' ability to identify their emotions. Emotional regulation acts like a balancing thermostat (18). Since the impact of improved emotional regulation processes on psychological issues in patients with myocardial infarction has been less studied, the present study examined the effectiveness of this type of psychotherapy on reducing depression, anxiety and stress in patients with myocardial infarction.

## Methods

The present study is a single case experimental in terms of data collection method in which the participants are evaluated in the baseline and treatment stages, respectively. The main objective of single-case designs in clinical work (trial) is to assist clinical professionals in evaluating the effect of a clinical intervention in changing the clients' behavior. The underlying logic of single-case design is similar to group design and the effect of intervention is examined by comparing the different conditions presented to the subject.

The performance of the subject in the pre-intervention or baseline stage is used to predict the subject's behavior in the future. The basic logic of single-case designs is similar to group design, and the effect of intervention is examined by comparing the different conditions presented to the subject, as it measures the impact of method in question on the subjects individually.. In the single-case experimental design, the control group does not exist and the baseline of each patient is considered as its control group (20). The statistical population of the present study was patients with myocardial infarction referring to Ayatollah Taleghani Hospital. In this study, convenience sampling method was used. 5 patients with myocardial infarction referred to Taleghani Hospital who had inclusion criteria were selected to participate in the treatment. The inclusion criteria were included the diagnosis of myocardial infarction by the cardiologist, the ability to attend sessions, the absence of severe mental illness (psychotic disorders, bipolar disorder, etc.), having at least the literacy of reading and writing. Exclusion criteria were included the absence of more than 2 sessions. Before the intervention, a pretest was used to determine the baseline, and the patient's condition was evaluated in terms of the variables. Then, the intervention was presented to participants in 8 sessions of 45 minutes, and they were evaluated at the end of each session. Prior to the start of the study, enough information on intervention was provided to the participants and the written informed consent was received from them to enter this research. Also, the appropriateness of the treatment for the participants was ensured. In order to maintain the principle of confidentiality, the information obtained from the evaluation sessions and questionnaires were encoded in such a way that the identity of the subjects were reserved and were only available to the researcher.

## Materials

Demographic characteristics were collected through personal information questionnaire that included age, sex, education level, marital status, history of referral to a psychologist or psychiatrist, history of smoking, alcohol and drugs.

## **Short form of depression, anxiety and stress scale (DASS-21):**

This scale was developed by Loeffond and Lovibond (1995) to simultaneously measure the severity of depression, anxiety and stress in individuals. The short form of the DASS-21 scale consists of 21 phrases related to the symptoms of negative emotions (depression, anxiety and stress). After reading each item, the subject graded the severity/frequency of the symptom in the questions during the last week using a 4-degree scale (between zero and three). Each of the three scales of depression, anxiety and stress has 7 questions, and the score is obtained through the total score of the questions related to it. Since the shortened form of the original scale consists of 42 questions, the final score of each subscale has to be doubled. Various studies have shown that the DASS-21 subscales have desirable psychometric properties (21). This scale has been studied by Samani and Jokar (2007), which have reported test re-test reliability for depression, anxiety and stress scales of 81, 76, 77, respectively and Cronbach's alpha for scales depression, anxiety, and stress have been reported 81, 74, 78, respectively (22).

The protocol used in this study was based on Grass's emotional regulation. The content of Grass's emotional regulation plan is summarized briefly. Session 1: Establishing a therapeutic relationship (rapport), introducing emotional regulation therapy, familiarizing with the subject of research and introductory explanations, expressing the main and the secondary goals, expressing the framework and rules, presenting the assignment, and feedback. Session 2: discussion about the nature of emotion and various kinds of it, a daily note card to record emotion, a practical exercise of emotion (emotional state experience), and a distinction between physical, mental and sensational dimensions of each emotion, assigning homework and feedback. Session 3: discussion about the emotion function in the process of adjustment and the role of emotion in establishing communication, how to organize and stimulate human behavior, being in an emotional state, completing self-assessment forms of emotional responses, completing the form of emotional vulnerability, grading, talk (discussion) about the context and cause of self vulnerability and identifying self-

regulation strategies: Complete the form of assessment of the patient's response to emotion (emotional skill and its effectiveness), assignment and feedback. Session 4: preventing social isolation and avoidance, providing a list of individual and social goals, identifying the situations that need to be restored, improved or changed, training problem-solving skills and interpersonal skills, assignment and feedback. Session 5: introducing James Grass's emotion regulation process model, introducing attention change skills, training of the skill to stop rumination and worry, attention skill training, assigning homework and feedback. Session 6: Discussing the role of the mind in producing, maintaining, increasing and decreasing emotional response, identifying false evaluations and its effects on emotional states, completing a list of ideas and evaluating false impressions (evaluations) on emotion, teaching re-evaluation strategies and modifying incorrect evaluation (misinterpretation), and as a result of relieving and adjusting negative emotions, giving a homework and feedback. Session 7: exposure; creating emotional states in session; confronting with emotion and avoiding avoidance; describing the physical, mental and sensational effects of emotion, in writing and verbal ways, evaluating (identifying the extent and the use of an avoidance strategy) and examining its emotional consequences. Skill learning as follows:

- Emotion expression
- Behavioral modification through changing the environmental reinforcing factors
- Emotional drainage strategy (abreaction)
- Changing the emotional outcomes through reverse operation
- Changing

- Creating physical relaxation through relaxation training
- Assignment
- Feedback

Session 8: Reassessment and planning for the application of training, assessing the reparation of a list of Individual goals, grading the patient's achievement, encouraging patient success, applying the proposed practical plan for the application of actual skills, program setting, assignment and Feedback (23).

### Results

A graphical or visual analysis was used to analyze the data. Based on the profile process, the effects of the independent variable on the dependent variable were examined. On the basis of dispersion (descent and ascension), scores, percentages of recovery, the permanent change index (RCI) and the effect size - firstly introduced by Jacobson and Trax (1991), were used. In this study, to calculate the recovery rate, the pre-test score is reduced from the post-test score and the principle is divided into the pre-test score, if the recovery rate is less than 50, then the results can be considered clinically significant (24).

The permanent change index indicates that the statistical change is due to the treatment intervention. For RCI to be clinically significant, the absolute value of the result should be equal to or greater than 1.96, which indicates that the results are mostly due to active factors and manipulation of the experimenter, rather than the measurement error (25).

$$RCI = \frac{X_{post} - X_{pre}}{\sqrt{2SE^2}}$$

**The descriptive statistics of participants in the demographic variables of employment, education, and marital status are presented in Table 1.**


**Table 1** Patients' descriptive findings

fifth	fourth	third	second	First	variables
۵۹	۷۰	۵۸	۵۹	۵۰	Age
Female	man	Female	man	man	Sex
guidance	Bachelor	guidance	guidance	Diploma	education
Married	Married	widow	Married	Married	marital status
-	-	-	-	-	History of referral to psychiatrist or psychologist
-	Alcohol sometimes	-	-	-	Drug use history Drug narcotic drugs

Recovery (improvement) and the effect of treatment in the subscales of depression, anxiety and stress in each subject

**Table 2** Comparison of depression scores of subjects and their recovery rates

RCI	Overall improvement	The recovery rate	Difference before and post test ازمون پیش و پس	Session 7	Session 6	Session 5	Session 4	Session 3	Session 2	Session 1	Basic line	Subjects
5/34		%62	۱۰	۶	۶	۸	۱۲	۱۴	۱۴	۱۸	۱۶	First
8/55	%68	%100	۱۶	۰	۲	۶	۱۰	۱۶	۱۲	۱۶	۱۶	second
7/48		%58	۱۴	۱۰	۱۲	۱۲	۱۸	۱۶	۲۲	۲۲	۲۴	third
6/41		%66	۱۲	۶	۱۰	۱۰	۱۴	۲۰	۲۲	۱۸	۱۸	fourth
5/88		%55	۱۱	9	۱۲	۱۲	۱۴	۱۶	۲۰	۱۸	۲۰	fifth


**Chart 1** Patients' depression score during eight sessions of therapy

The RCI score for all five subjects is greater than 1.96, meaning that the results are mostly due to active factors and manipulation of the

experimenter, rather than the measurement error. The percentage of recovery for each of the five subjects is remarkable (considerable), and in general, it can be said that in all five subjects, the overall recovery index is 68%, which is significant. Therefore, based on these results, it can be concluded that emotional regulation therapy has improved depression in patients with myocardial infarction in the post-intervention phase than before intervention.

**Table 2** Comparison of anxiety scores of subjects and their recovery rates


RCI	Overall improvement	The recovery rate	Treatment sessions									Subjects
			Difference before and post test	Session 7	Session 6	Session 5	Session 4	Session 3	Session 2	Session 1	Basic line	
5/23	%69.8	%71	۱۰	۴	۸	۸	۶	۱۰	۱۲	۱۲	۱۴	First
10/47		%90	۲۰	۲	۴	۸	۱۰	۱۶	۱۴	۱۸	۲۲	second
7/32		%54	۸۱	8	۸	۱۰	12	10	۱۸	۲۰	۲۲	third
5/23		%65	۶۲	۶	۱۰	۱۰	۱۴	۱۶	۲۰	۱۸	۱۶	fourth
4/18		%57	۸	۶	۸	۶	۱۰	۱۴	۱۴	۱۶	۱۴	fifth

**Chart 2** Patients' anxiety score during eight sessions of therapy

The RCI score for all five subjects is greater than 1.96, meaning that the results are mostly due to active factors and manipulation of the experimenter, rather than the measurement error. The percentage of recovery for each of the five subjects is remarkable (considerable), and in general, it can be said that in all five subjects, the overall recovery index is 69.8% which is significant. Therefore, based on these results, it can be concluded that emotional regulation therapy has improved anxiety in patients with myocardial infarction in the post-intervention phase than before intervention.

**Table 3** Comparison of stress scores of subjects and their recovery rates

RCI	Overall improvement	The recovery rate	Treatment sessions									Subjects
			Difference before and post test	Session 7	Session 6	Session 5	Session 4	Session 3	Session 2	Session 1	Basic line	
3/42	%59	%50	۱۰	۱۰	۱۲	۱۶	۱۸	۲۰	۲۲	۲۸	۲۰	First
8/90		%76	۲۶	۸	۱۲	۱۶	۲۴	۳۰	۳۴	۳۲	۳۴	second
8/21		%70	۲۴	۱۰	۱۰	۱۶	۲۲	۲۸	۲۸	۳۲	۳۴	third
4/79		%53	۱۴	12	۱۴	۱۶	۲۰	۱۸	۲۲	۲۴	۲۶	fourth
3/74		%45	۱۰	12	۱۶	۱۴	۱۸	۲۰	۱۸	۲۰	۲۲	fifth



**Chart 3** Patients' stress score during eight sessions of therapy

The RCI score for all five subjects is greater than 1.96, meaning that the results are mostly due to active factors and manipulation of the experimenter, rather than the measurement error. The percentage of recovery for each of the five subjects is remarkable (considerable), and in general, it can be said that in all five subjects, the overall recovery index is %59, which is significant. Therefore, based on these results, it can be concluded that emotional regulation therapy has improved stress in patients with myocardial infarction in the post-intervention phase than before intervention.

### Discussion

The purpose of this study was to determine the effectiveness of emotional regulation therapy on the improvement of depression, anxiety and stress in patients with myocardial infarction. In this study, increasing the rate of recovery rate, high effect size, and reviewing the process of change showed that the intervention of emotional-cognitive regulation therapy was effective on depression, anxiety and stress, and clinically resulted in significant changes before and after intervention.

The results of current study are consistent with the findings of the studies conducted by Martin and Dahlon 2005 (26), Schwarz et al. 2008 (27), Linde Cobst in 2009 (28), Quid et al (2010) (29), Abdi et al., 2010(30), Manin and Fresco 2013 (31), Sobhi et al., In 2015 (32), Foroughy in 2016 (33), Shast Fooladi and Monsheyee, 2016 (34).

In the explanation of the effectiveness of emotional regulation therapy on depression, anxiety and stress the studies have shown

emotional disorders can be conceptualized through inappropriate use of emotional regulation strategies. The emotional regulation processes are damaged in patients with depression, and depressed people have a lower capacity to tolerate negative emotions and little awareness of their emotions compared to normal people. They use dysfunctional coping skills to cope with their emotions (35); and use maladaptive strategies such as rumination, self-blame, other-blame and catastrophizing. Studies have shown significant positive correlation between depression (36), anxiety, stress (37), and dysfunctional components of cognitive-emotional regulation, and suggested that these symptoms as the most important risk factors for cardiovascular disease (38). In this regard, the study by Schroevers et al. showed that patients who were using adaptive strategies in dealing with stressful events had less depression and had higher levels of health and well-being (27).

According to the Grass's model, the emotional response takes a path that can be examined at several points (dealing with position, attention, evaluation, and response); therefore, emotional regulation can occur in any of the points of this path. Emotional regulation therapy causes improved emotional regulation patterns, increased individual's emotional awareness, being adapted to stressful events of life, decreased rumination, positive thinking and changes individual assessments (evaluations) (39) and helps individuals to manage their emotions (39, 40).

### Conflict of interests

Authors declare no conflict of interests.

### References

1. Kasper Hauser J f, Longo, Lvskalzv Translation: khodaei, Khamene. Harrison's Principles of Internal Medicine. arjmand Tehran publisher. 2015.
2. Vahidian Azimi A AF AF, Kazemnezhad A. Assesing risk factors coronary artery disease. 3th Iran-Arab (Middle East) Cardiovascular Congress 2009.
3. Satianda s. Treatment of heart disease. Translation Mousavi nasab jFETfp.
4. Sullivan MJ WL, Terry J, Brantley J, Charles A, McGee V, et al. The Support, Education, and Research in Chronic Heart Failure Study (SEARCH): a mindfulness-



based psychoeducational intervention improves depression and clinical symptoms in patients with chronic heart failure. *American heart journal*. 2009;157(1):84-90.

5. Corvera-Tindel T DL, Aquino C, Roper J, Dracup K. The role of physical functioning and depression in quality of life in patients with heart failure. *Journal of Cardiac Failure*. 2003;9(5):S5.

6. Riedinger MS DK, Brecht M-L, Investigators S. Quality of life in women with heart failure, normative groups, and patients with other chronic conditions. *American Journal of Critical Care*. 2002;11(3):211-9.

7. Meijer A CH BE, Thombs BD, van Melle JP, de Jonge P. Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis of 25 years of research. *General hospital psychiatry* 2011; 33(3): 203-16.

8. Behnammoghadam M BA STEoEMDaREodipwMIMia-mfuJ.

9. Lane D CD, Ring C, Beevers DG, Lip GY. The prevalence and persistence of depression and anxiety following myocardial infarction. *British journal of health psychology*. 2002;7(1):11-21.

10. Parakh K TB, Fauerbach JA, Bush DE, Ziegelstein RC. Effect of depression on late (8 years) mortality after myocardial infarction. *The American journal of cardiology*. 2008;101(5):602-6.

11. Frasure-Smith N LFDacadH-.

12. Hanssen TA NJ, Eide GE, Bjelland I, Rokne B. Anxiety and depression after acute myocardial infarction: an 18-month follow-up study with repeated measures and comparison with a reference population. *European Journal of Cardiovascular Prevention & Rehabilitation*. 2009;16(6):651-9.

13. Dunderdale K TD, Miles JN, Beer SF, Furze G. Quality-of-life measurement in chronic heart failure: do we take account of the patient perspective? *European journal of heart failure*. 2005;7(4):572-82.

14. Kaplan Sadock SoPBSCPVITR HS, Khosro. First Edition. Tehran: Arjmand 1387.

15. keynoosh h PoCR, (First Edition . Tehran faravan publications), 1385. .

16. Arora D AM, Katyal V, Anand V. Anxiety and well-being among acute coronary syndrome patients: overtime. *J Indian Acad Appl Psychol*. 2010;36:79-88.

17. Rahimian b, Sadat alami. 6-4 months post-MI patients change health behaviors. *Iran Journal of Nursing*. 2000;13 (24): 8-15.

18. Napolitano. *Emotion Regulation in Psychotherapy techniques Practical Guide*. 1392.

19.

Shaughnessy, J.J., Zechmeister. E.B, Research, methods in Psychology. New York: McGrawHill. 1997.

20. Anderson CM, Kim C. Evaluating treatment efficacy with single-case designs. Roberts MC, Ilardi SS, editors, *Handbook of research methods in clinical psychology*. Malden, MA: Blackwell; 2003;73-91.

21. Lovibond SM, Lovibond PFM. *Manual for the Depression Anxiety Stress Scales*. 2nd ed. Australia, Sydney: Psychology Formulation; 1995.

22. Samani S Joukar B, . A Study The Reliability And Validity Of The Short Form Of The Depression Anxiety Stress Scale (DASS-21). *Social Sciences And Humanities Of Shiraz University* , Volume 26 , Number 3 (52) (Special Issue In Education). Fall 2007; 65 -77

23. Salehi A, Baghban I, Bahrami F, Ahmadi SA. The effect of emotion regulation training based on dialectical behavior therapy and gross process model on symptoms of emotional problems. *Zahedan Journal of Research in Medical Sciences*. 2012;14(2):49-55.

24. Soliemanian AA, Naghinasab Ardehaee F and S. "The Effectiveness of Systemic Sex Therapy on Sexual Desire Improvement in Women With Hypoactive Sexual Desire Disorder(HSDD). " *Iranian Journal of Psychiatric Nursing (IJPNA)* A (2016) ;3(4): 60-70.

25. Naseri A, F. Sohrabi A. Bardhaili and M Philosophical race "The effectiveness of a two-dimensional plan therapy in the treatment of heroin addiction associated with antisocial personality disorder. " *Clinical Psychology Studies* 6(2015); (18): 75-97.

26. Martin RC, Dahlen ER. Cognitive emotion regulation in the prediction of depression, anxiety, stress, and anger. *Personality and Individual Differences*. 2005;39(7):1249-60.

27. Schroevers M, Kraaij V, Garnefski N. How do cancer patients manage unattainable personal goals and regulate their emotions.



British journal of health psychology. 2008;13(3):551-62.

28. Lindquist R, Windenburg D, Savik K, Bronas U. Pilot of Stress Reduction Strategies for Patients After a Coronary Event. *Cleveland Clinic Journal of Medicine*. 2009;76(Suppl 2):S97a-Sa.

29. Quoidbach J, Berry EV, Hansenne M, Mikolajczak M. Positive emotion regulation and well-being: Comparing the impact of eight savoring and dampening strategies. *Personality and Individual Differences*. 2010;49(5):368-73.

30. Abdi S, Babapoor J, Fathi H. Relationship between Cognitive Emotion Regulation Styles and General Health among University Students. *Military Medical University, Iran*. 2010.

31. Mennin D, Fresco DM., Heimberg RG, Ciesla J. Randomized control trial of emotion regulation therapy for generalized anxiety disorder and comorbid depression. In M. Fraire & T. Ollendick (Chairs), *Emotion regulation in anxiety disorders across the lifespan: Comorbidity and treatment*. Symposium presented at the annual meeting of the Anxiety Disorders Association of America, Arlington., 2012.

32. Sobhi-Gharamaleki N PP, Aghajani S, Narimani M. Effectiveness of Emotion Regulation Training on Reduction of Anxiety, Stress and Depression Symptoms among University Students. *Iranian Journal of Health Education and Health Promotion*. 2015;3(1):5-13.

33. Foroughi A. Examination fitness of Emotion Dysregulation Model and comparison efficacy of Emotion Regulation Therapy with Integrative Psychotherapy in Generalized Anxiety Disorder (GAD): symptom reduction, improving function, enhancement of emotion regulation skills and decrease interpersonal problems. 2016.

34. Shastfooladi M, Manshaei G. The Effectiveness of Emotion Regulation Group Training on Perceived Stress in Women with Breast Cancer in Isfahan. 2016.

35. Kashdan TB BV, Forsyth JP, Steger MF. Experiential avoidance as a generalized psychological vulnerability: comparisons with coping and emotion regulation strategies. *Behav Res Ther* 2006; 44(9): 1301-20.

36. Garnefski N, Kraaij V. Relationships between cognitive emotion regulation

strategies and depressive symptoms: A comparative study of five specific samples. *Personality and Individual Differences*. 2006;40(8):1659-69.

37. Watson D, & Clark, L. A. Negative affectivity the disposition to experience aversive emotion states. *Psychological Bulletin*, 96, 465-490. 1984.

38. Kubzansky LD, Kawachi I. Going to the heart of the matter: do negative emotions cause coronary heart disease? *Journal of psychosomatic research*. 2000;48(4):323-37.

39. Gross JJ, Thompson RA. *Emotion regulation: Conceptual foundations*. 2007.

40. Ehring T FS, Schnulle J, Bosterling A, Tuschen-Caffier B. Characteristics of emotion regulation in recovered depressed versus never depressed individuals. *Personality and Individual Differences* 2008; 44(7): 1574-84.