

// ( )  
//  
//

\*

---

paired t-test  
/ ± / ( % % ) :  
/ ± / / ± / FEV0.5/FVC ± / / ± / FEV1/FVC :  
FEV1/PEFR (P < / ) / ± / ± / MEF50 / ± / / ± / FEV25-75

---

(.) Adenotonsillar- (ATH) Hypertrophy  
( )  
ATH (.)  
Obstructive Sleep Apnea Syndrome(OSAS)  
%  
ATH (.) ( )  
ATH (.)  
( )

---

PT Clotting Time Bleeding Time

PTT

Yadav

.( )

/ Maurizi

Minato

.( )

(Forced Expiratory Volume in First Second )

FVC (Functional Vital Capacity) FEV1

FEV0.5 (FEV1/FVC) FVC FEV1

( )

(Forced Expiratory Volume in half a second)

FEV25-75 (FEV0.5/FVC) FVC FEV0.5

(Forced Expiratory Volume 25%-75%)

MEF50 (Peak Expiratory Flow Rate) PEFR

)

(Maximum Expiratory Flow at 50% of vital capacity)

)

(( ) Brodsky

FEV1/ PEFR

.(

## Paired t-test

P- Value				
/	+ /	± /	/ ± /	<b>FEV1/FVC</b> ( )
/	+ /	/ ± /	/ ± /	<b>FEV0.5/FVC</b> ( )
/	+ /	/ ± /	/ ± /	<b>FEV25-75</b> ( )
/	+	/ ± /	± /	<b>MEF50</b> ( )
/	/	/ ± /	/ ± /	<b>FEV1/FEV0.5</b> ( )
/	/	/ ± /	/ ± /	<b>FEV1/PEFR</b> ( )

(% ) )

( ( % )

( ) / ± /

+  
(%)

( / %)

( / %)

FEV0.5/FVC

.( )

( )

FEV1/PEFR

.(P &lt; / )

MEF50

(% / )

FEV0.5/FVC

.(P= / )

( ) % / (Flow Volume Plot)  
% (Forced Inspiratory Flow)

Kavukcu

%

%

%

(Peak Expiratory Flow Rate)

(FEV1/FEV0.5) /

(.)

( )

Yadav

/

FEV1/PEFR FEV1/FEV0.5

FEF% 50/FIF% 50

(.)

FEV1

(.)

FEV1

FEV1

(FEV1/ PEFR)

) FVC

( ) FEV1 ( PEFR (P= / )

FEV0.5 (FEV1/FVC) FVC FEV1

FEV1

FEV1/FVC ( / )

FEV1/FEV0.5

( ) FEV25-75 ( )

( ) MEF50 /

FEV1/FEV0.5

FEV1/FEV0.5

(P < / )

Maurizi

---

## REFERENCES

---

1. Greenfeld M, Tauman R, DeRowe A, Sivan Y. Obstructive sleep apnea syndrome due to adenotonsillar hypertrophy in infants. *Int J Pediatr Otorhinolaryngol* 2003; 67(10): 1055-60.
2. Tezer MS, Karanfil A, Aktas D. Association between adenoidal-nasopharyngeal ratio and right ventricular diastolic functions in children with adenoid hypertrophy causing upper airway obstruction. *Int J Pediatr Otorhinolaryngol* 2005; 69(9): 1169-73.
3. Mora R, Salami A, Passali FM, Mora F, Cordone MP, Ottoboni S, et al. OSAS in children. *Int J Pediatr Otorhinolaryngol* 2003; 67 Suppl 1: 229-31.
4. Kurnatowski P, Putynski L, Lapienis M, Kowalska B. Neurocognitive abilities in children with adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol* 2006; 70(3): 419-24.
5. Yadav SP, Dodeja OP, Gupta KB, Chanda R. Pulmonary function tests in children with adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol* 2003; 67(2): 121-5.
6. Ersoy B, Yuceturk AV, Taneli F, Urk V, Uyanik BS. Changes in growth pattern, body composition and biochemical markers of growth after adenotonsillectomy in prepubertal children. *Int J Pediatr Otorhinolaryngol* 2005; 69: 1175-81.
7. Gorur K, Doven O, Unal M, Akkus N, Ozcan C. Preoperative and postoperative cardiac and clinical findings of patients with adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol* 2001; 59(1):41-6.
8. Pac A, Karadag A, Kurtaran H, Aktas D. Comparison of cardiac function and valvular damage in children with and without adenotonsillar hypertrophy. *Int J Pediatr Otorhinolaryngol* 2005; 69(4): 527-32.
9. Kharb S, Yadav SP, Singh H, Singh GP. Effect of adenotonsillectomy on arterial blood gases and acid-base balance. *Int. J. Pediatr Otorhinolaryngol* 1998; 43: 213-15.
10. Maurizi M, Paludetti G, Todisco T, Dottorini M, Grassi V. Pulmonary function studies in adenoid hypertrophy (abstract). *Int J Pediatr Otorhinolaryngol* 1980; 2(3): 243-50.

- 
11. Yilmaz MD, Onrat E, Altuntas A, Kaya D, Kahveci OK, Ozel O, et al. The effects of tonsillectomy and adenoidectomy on pulmonary arterial pressure in children. *Am J Otolaryngol* 2005; 26(1): 18-21.
  12. Kavukcu S, Coskun S, Cevik N , Kuscu B , Akkoclu A. The importance of pulmonary function tests in adenotonsillectomy indications (abstract). *Indian J Pediatr* 1993; 60(2): 249-55.