

//
//

()

*

()

:

(Cross-sectional)

:

DMSA

()

ESR

%

OR

Chi-square

:

(% /)

($P < /$)

ESR

DMSA

:

DMSA

:

() %

DMSA

()

()

%
 (% /)
 (Many)
 % /
 (%)
 ESR

(Mild- Moderate)

(Cross-sectional)

% /
 %
 % / %
 (% /) (% /)

/ °C
 (WBC/HPF)

VCUG

DMSA

()

ESR

(% /)

(% /)

ESR

ESR ()

(P< /)

VCUG

/

%

/ /

(% /)

ESR

%

(CI)

ESR

(% /)

(OR= / P< /)

Chi-square

% /

% /

(OR) Odd's Ratio

(OR= / P< /)

% /

%

(OR= / P< /)

DMSA

		OR	P-value	
	(.)			
	Ditchfield	—	P< /	(/) (/)
		/	P< /	(/) (/)
		/	P< /	() ()
	(P< /)	—	P< /	(/) (/)
	(. ())	/	P< /	(/) (/)
				(Many)
	DMSA		P< /	(/) (/)
CPR	Lai	/	P< /	(/) (/)
	Positive Predictive Value			
	LEE			(.)
	CPR			
	(. ())			
	CRP Ghio			()
ESR	(. ())			
				% /
	% / (/)			
	(% /)			
				(P< /)
				()

() % % /

% /

ESR DMSA (Mild- Moderate)
 () %
 ()
 % /

()

) %

(/) % (

(OR = / P < /)

()

% /

REFERENCES

1. Behrman R E, Kliegman R M, Jenson H B. Nelson Textbook of Pediatrics. Philadelphia: Saunders. 2004; P: 1789.
2. Jakobsson B, Tolstedt L, Svensson L, Soderlundh S, Berg U. 99m Technetium dimercaptosuccinic acid scan in the diagnosis of acute pyelonephritis in children: relation to clinical and radiological findings. *Pediatr Nephrol* 1993; 7(5): 591-2.
3. Biggie A, Dardanelli LN, Cussant P, et al. Prognostic value of the acute DMSA scan in children with first urinary tract infection. *Pediatr Nephrol* 2001; 16: 800-4.
4. Nikbakhsh A A, Yekta Z, Mahmoodzadeh H, Karamiyar M, Fazel M. Technetium Tc 99m Dimercaptosuccinic Acid Renal Scintigraphy in Diagnosis of Urinary Tract Infections in Children with Negative Culture. *Urol J* 2006; 3: 139-42.
5. Christian MT, McColl JH, MacKenzie JR, Beattie TJ. Risk assessment of renal cortical scarring with urinary tract infection by clinical features and ultrasonography. *Arch Dis Child* 2000; 82: 376-80.

-
6. Michael A, Ditchfield J, de Campo F, Terrance M, Notan D, Keith Grimwood K, et al. Risk Factors in the Development of Early Renal Cortical Defects in Children with Urinary Tract infection. *AJR* 1994; 162: 1393-97.
 7. Levtchenko EN, Lahy C, Levy J, Ham HR, Piepsz A. Role of Tc-99m DMSA scintigraphy in the diagnosis of culture negative pyelonephritis. *Pediatr Nephrol*. 2001; 16: 503-6.
 8. Nammalwar BR, Vijayakumar M, Sankar J, Ramnath B, Prahlad N. Evaluation of the use of DMSA in culture positive UTI and culture negative acute pyelonephritis. *Indian Pediatr*. 2005; 42: 691-6.
 9. Lai SW, Ng KC. Retrospective analysis of inflammatory parameters in acute pyelonephritis. *Scand J Urol Nephrol* 2003; 37(3): 250-2.
 10. Lee BF, Chiou YY, Chuang CM, Wu PS, Wu YC, Chiu NT. Evolution of differential renal function after acute pyelonephritis. *Nucl Med Commun* 2002; 23: 1005-8.
 11. Ghireo L, Cracco AT, Sartor M, Comacchio S, Zacchello G, Dall'Amico R, et al. Retrospective study of children with acute pyelonephritis, evaluation of bacterial etiology, antimicrobial susceptibility, drug management and imaging studies. *Nephron* 2002; 90(1): 8-15.
 12. Ditchfield MR, Nadel HR. The DMSA scan in pediatric urinary tract infection. *Australas Radiol* 1998; 42 (4): 318-20.
 13. Ataei N, Madani A, Habibi R, Khorasani M. Evaluation of acute pyelonephritis with DMSA scans in children presenting after the age of 5 years. *Pediatr Nephrol* 2005; 20: 1439-44.
 14. Moorthy I, Easty M, McHugh K, Ridout D, Biassoni L, Gordon I. The presence of vesicoureteric reflux does not identify a population at risk for renal scarring following a first urinary tract infection. *Arch Dis Child* 2005; 90: 733-6.
 15. Lee JH, Son CH, Lee MS, Park YS. Vesicoureteral reflux increases the risk of renal scars: a study of unilateral reflux. *Pediatr Nephrol* 2006 Sep; 21(9): 1281-4.
 16. Imperiale A, Olianti C, Sestini S, Materassi M, Seracini D, Ienuso R, et al. 123I-hippuran renal scintigraphy with evaluation of single-kidney clearance for predicting renal scarring after acute urinary tract infection: comparison with (99m) Tc-DMSA scanning. *J Nucl Med* 2003; 44 (11): 1755-60.