

// ()
// , ,

*

(Urinary Tract Infection) UTI :

(Macrophage Migration Inhibitory Factor) MIF

MIF

MIF

)

:

midstream

(

ELISA

MIF

One Way ANOVA Independent-t test

SPSS 13

(Receiver Operating Curve) ROC

(SEM = /) pg/ μ mol creatinine (A)

MIF/Cr

:

(C) / (SEM = /) pg/ μ mol creatinine (B) /

(P < /) B A / (SEM = /) pg/ μ mol creatinine

MIF/Cr

ROC

(P < /) C

[Area Under Curve(AUC)= / P < /]

/ pg/ μ mol creatinine

(AUC= / P < /)

/ pg/ μ mol creatinine

MIF

:

MIF/Cr

MIF/Cr

MIF/Cr

:

(Urinary Tract Infection)

UTI

()

(Convenience)

()

MIF

°C

midstream

MIF

(Macrophage Migration Inhibitory Factor)

×g

T

ELISA

MIF

MIF

(PMN)

()

ESR

DMSA

MIF

)

()

(

MIF

SPSS.13

MIF

SEM (SD)
(Standard Error of Mean)
MIF
Mann-Whitney U-test
Independent-t test
(A)
(C) (B)
B A
(SEM = /) / (SEM = /) C AUC
(SEM = /) / (Area Under Curve)
ANCOVA

MIF/Cr		Cr MIF	
MIF/Cr	Cr	MIF	
(pg/ μ mol)	(mg/dl)	(pg)	()
/	/	/	(%) :
(SEM= /)	(SEM= /)	(SEM= /)	(SEM= /) (%) :
/	/	/	(%) :
(SEM= /)	(SEM= /)	(SEM= /)	(SEM= /) (%) :
/	/	/	(% /) :
(SEM= /)	(SEM= /)	(SEM= /)	(SEM= /) (% /) :


MIF
MIF/Cr
B A
pg/ μ mol creatinine :
MIF/Cr / (SEM = /)
ESR () / (SEM = /) pg/ μ mol creatinine (A)
(P < /) MIF (P= /) pg/ μ mol (B)
(P= /) PMN .C / (SEM = /) creatinine
B A MIF/Cr
(p < /) C (p < /)

B A

ANOVA

(P= /)

MIF/Cr



MIF/Cr

MIF/Cr

(P< /)

(P< /)

(P= /)

MIF/Cr

B A

P-value	B	A	
< / *	(SEM= /)	(SEM= /)	(pg) MIF
/ *	(SEM= /)	(SEM= /)	(mg/dl) Cr
< / *	(SEM= /)	(SEM= /)	(pg/μmol) MIF/Cr
/	(SEM= /)	(SEM= /)	()
/	(SEM= /)	(SEM= /)	(/μl) WBC
/	(SEM=)	(SEM= /)	(/μl) PMN
/ *	(SEM= /)	(SEM= /)	(mm/h) ESR

(P< /)

*

% / %
/ pg/ μ mol creatinine
% %

ROC

MIF/Cr


(AUC= / P < /)
(AUC= / P < /)

MIF/Cr

ROC

/ pg/ μ mol creatinine (cut-point)/ pg/ μ mol creatinine

(AUC)


AUC_{MIF/Cr} = / (P= /)](P= /) AUC_{PMN} = / (P= /).[AUC_{WBC} = / (P= /) AUC_{ESR} = // pg/ μ mol creatinine .

(Receiver Operating Curve) ROC

(pg/ μ mol) MIF/Cr

= /]

[P < / (AUC)





(Receiver Operating Curve) ROC

(pg/ μ mol) MIF/Cr

= /]

[P < / (AUC)




(Receiver (AUC) Operating Curve) ROC

MIF/Cr (P = / AUC = /)
 ESR (P = / AUC = /)
 WBC (P = / AUC = /)
 = /) Cr (P = / AUC = /)
 (P = / AUC

) MIF/Cr (pg/μmol)
 MIF/Cr / pg/μmol
 = %)
 . (= % /

(MIF)
 () T
 MIF
 ()
 MIF
 ()
 MIF



) MIF/Cr (pg/μmol)
 / pg/μmol MIF/Cr
 (= % = %)

/

MIF

MIF

MIF/Cr

Meyer-Siegler

MIF

MIF/Cr

.()

MIF/Cr

.()

Meyer-Siegler

pg/ μ mol creatinine / pg/ μ mol creatinine

MIF ELISA

/

MIF

.()

MIF/Cr

.()

MIF

REFERENCES

1. Cunningham RJ, Carvajal HF, Passey RB. Urinary LDH iso-enzyme 5 excretion in experimental pyelonephritis. Br J Exp Pathol 1977; 58: 220.

2. Carvajal HF, Passey RB, Berger M, Travis LB, Lorentz WB. Urinary lactic dehydrogenase isoenzyme 5 in the differential diagnosis of kidney and bladder infections. *Kidney International* 1975; 8: 176-84.
3. Morand EF, Bucala R, Leech M. Macrophage migration inhibitory factor: an emerging therapeutic target in rheumatoid arthritis. *Arthritis Rheum* 2003; 48: 291-9.
4. Bacher M, Metz CN, Calandra T, Mayer K, Chesney J, Lohoff M et al. An essential regulatory role for macrophage migration inhibitory factor in T-cell activation. *Proc Natl Acad Sci* 1996; 93: 7849-54.
5. Metz CN, Bucala R. Role of macrophage migration inhibitory factor in the regulation of the immune response. *Adv Immunol* 1997; 66: 197-223.
6. Metz CN, Bucala R. Cytokine reference: a compendium of cytokines and other mediators of host defense. Vol: 1. San Diego: Academic Press. 2001; PP: 703-16.
7. Huang XR, Hui CWC, Chen Y-X, Chun B, Wong Y, Fung PCW, et al. Macrophage migration inhibitory factor is an important mediator in the pathogenesis of gastric inflammation in rats. *Gastroenterol* 2001; 121: 619-30.
8. Ohkawara T, Nishihira J, Takeda J, Hige S, Kato M, Sugiyama T, et al. Amelioration of dextran sulfate sodium-induced colitis by anti-macrophage migration inhibitory factor antibody in mice. *Gastroenterol* 2002; 123: 256-70.
9. Bacher M, Meinhardt A, Lan HY, Dhabhar FS, Mu W, Metz CN, et al. MIF expression in the rat brain: implications for neuronal function. *Mol Medicine* 1998; 4: 217-30.
10. Vera PL, Iczkowski KA, Leng L, Bucala R, Meyer-Siegler KL. Macrophage migration inhibitory factor is released as a complex with alpha1-inhibitor-3 in the intraluminal fluid during bladder inflammation in the rat. *J Urol* 2005; 174: 338-43.
11. Meyer-Siegler KL, Iczkowski KA, Vera PL. Macrophage migration inhibitory factor is increased in the urine of patients with urinary tract infection: macrophage migration inhibitory factor-protein complexes in human urine. *J Urol* 2006; 175(4): 1523-8.
12. Vera PL, Meyer-Siegler KL. Anatomical location of Macrophage Migration Inhibitory Factor in urogenital tissues, peripheral ganglia and lumbosacral spinal cord of the rat. *BMC Neurosci* 2003; 4: 17.
13. Meyer-Siegler KL, Vera PL, Ordorica RC, Fernandez HL. Proinflammatory cytokine up-regulation in central and peripheral tissue innervating the bladder in an endotoxin induced model of rat bladder inflammation. *J Urol* 2005; 172: 1507-9.
14. Vera PL, Ordorica RC, Meyer-Siegler KL. Hydrochloric acid-induced changes in macrophage migration inhibitory factor in the bladder, peripheral and central nervous system of the rat. *J Urol* 2003; 170: 623-7.