

//
//

()

*



" "

()

Hannalla

($P < /$)

mohsen_rouzrokh@yahoo.com

() .()

.()

()

N₂O

.()

ACTH

/

.()

cc)

() .()

()

Diplus)

.(Radium

/

(Hannalla)

Points	0	1	2
Blood pressure	BP=10% preoperative	BP>20% preoperative	BP> 30% preoperative
Crying	Not crying	Crying or responding to TLC *	Crying but not responding to TLC *
Movement	None	Restless	Trashing
Agitation	Child asleep or calm	Mild	Hysterical
Posture	No special posture	Flexing leg and thigh	Holding groin

* TLC: Tender Loving Care

(P< /) (P< /)

(P= /)

(P< /) (P< /)

(P= /)

()

()

()

(P< /)

(P< /)

(P< /)

(P< /)

(P< /)

(P= /)

(P= /)

(P= /)

(P< /)

(P< /)

(Hannalla)

(P= /)

/ ± /

/ ± /

(P= /)

()

()

*

			Mean ± SD	Mean ± SD	Mean ± SD	
/	/	/	/ ± /	/ ± /	/ ± /	μU/ml
/	/	/	/ ± /	/ ± /	/ ± /	mg/dL
/	/	/	/ ± /	/ ± /	/ ± /	μg/dl
/	/	/	/ ± /	/ ± /	/ ± /	mIU/L


* Paired t test (P<0.05)

()


*

			Mean ± SD	Mean ± SD	Mean ± SD	
/	/	/	/ ± /	/ ± /	/ ± /	μU/ml
/	< /	< /	/ ± /	/ ± /	/ ± /	mg/dL
/	< /	< /	/ ± /	/ ± /	/ ± /	μg/dL
/	/	/	/ ± /	/ ± /	/ ± /	μg/dL
/	< /	/	/ ± /	/ ± /	/ ± /	μg/dL
< /	/	< /	/ ± /	/ ± /	/ ± /	mIU/L
< /	/	< /	/ ± /	/ ± /	/ ± /	mIU/L


* t-test (P<0.05)




(ng/ml :)



(> μg/IU/ml :)



(/ ng/ml :)



(mg/ml :)

()
()

ACTH

Castejan-Casadoy

ACTH

()

()

Anand ()

George Sakellaris

()

()

()

()

()

()

REFERENCES

1. Denman WT, Ballantyme J. The Massachusetts general Hospital Hand book of pain management. 2nd ed. 2006; part: 22.
2. MC Donald AJ. Cooper Mg: patient-controlled analgesia: An appropriate method of pain control in children. *Paediatr Drugs* 2001; 3: 273-84.
3. Conroy JM, Othersen HB Jr, Dorman BH, Gottesman JD, Wallace CT, Brahen NH. A comparison of wound instillation and caudal block for analgesia following pediatric inguinal hernioaphy. *J Pediatr Surg* 1993; 28; 565-667.
4. Anand KJS, Agnsley-Green A. Measuring the severity of surgical stress in newborn infants. *J Pediatr Surg* 1988; 23: 297-305.
5. Anand KJ, Brown MJ, Causon RC, Christofides ND, Bloom SR, Aynsley-Green A. Can the human neonate mount an endocrine and metabolic response to surgery? *J Pediatr Surg* 1985; 20:41-48
6. Anand KJS Hickey PR. Halothane-morphine compared with high-dose sufentanil for anesthesia and postoperative analgesia in neonatal cardiac surgery. *N Engl J Med* 1992; 326: 1-9.
7. Håkanson E, Rutberg H, Jorfeldt L, Wiklund L. Endocrine and metabolic responses after standardized moderate surgical trauma: Influence of age and sex. *Clin Physiol* 1984; 4: 461-73.
8. Ishida H, Murata N, Yamada H, Nomura T, Shimomura K, Fujioka M, et al. Effect of CO2 pneumoperitoneum on growth of liver micrometastases in a rabbit model. *World J Surg* 2000; 24: 1004-8.
9. Rutberg H, Hakanson E, Anderberg B. Effects of the extradural administration of morphine, or Bupivacaine. *Br J Anasth* 1984; 86: 611-8.
10. Thorell A, Efendic S, Gutniak M, Häggmark T, Ljungqvist O. Insulin resistance after abdominal support in pediatrics. *Nutrition* 1998; 14: 130-42.
11. Casey WF, Rice LJ, Hannallah RS, Broadman L, Norden JM, Guzzetta P. A comparison between bupivacaine instillation versus ilioinguinal/iliohypogastric nerve block for postoperative analgesia following inguinal hernioraphy in children. *Anesthesiology* 1990; 72: 637-9.
12. Ward Platt MP, Tarbit MJ, Aynsley-Green A. The effects of anesthesia and surgery on metabolic homeostasis in infancy and childhood. *J Pediatr Surg* 1990; 25: 472-8.
13. Reid MF, Harris R Phillips RD. Day-case herniotomy in children. A comparison of ilioinguinal nerve block and wound infiltration for postoperative analgesia. *Anaesthesia* 1987; 42: 658-61.

14. Anand KJS. The stress response to surgical trauma: Form physiological basis to therapeutic implications. *Prog Food Nutr Sci* 1986; 10: 67-132.
15. Solak M, Ulusiy H, Sarihan H. Effects of caudal block on cortisol and prolactin responses to postoperative pain in children. *Eur J Pediatr Surg* 2000; 10: 219-23.
16. Okur H, Hucukaydin M, Muhtaroglu S. Effects of bupivacaine on beta-endorphin and cortisol release and postoperative pain following inguinal herniorraphy in children. *Peadiatr Surg Int* 1995; 11: 41-4.
17. Castejón-Casado J, Moreno-Prieto M, Valladares-Mendías JC, Alaminos-Mingorance M, López-Candel E, Ramírez-Navarro A. Hormonal response to sargical stress in school children. *Eur J Pediatr Surg* 2001; 11: 44-7.
18. Reiz S, Haggmark S, Johansson G, Nath S. Cardiotoxicity of ropivacaine: A new amide local anesthesia agent. *Acta Anaesthesiol Scand* 1989; 33(2): 93-8.
19. Wolf AR, Eyres RL, Laussen PC, Edwards J, Stanley IJ, Rowe P, et al: Effect of extradural analgesia on stress response to abdominal surgery in infants. *Br J Anaesth* 1993; 70: 654-60.
20. akellaris G, Petrakis I, Makatounaki K, Arbiros I, Karkavitsas N, Charissis G. Effects of Ropivacaine Infiltration on cortisol and prolactin responses to postoperative pain after inguinal hernioraphy in children. *J Pediatr Surg* 2004; 39(9): 1400-3.
21. Luz G, Innerhofer P, Haussler B, Oswald E, Salner E, Sparr H. Comparison of ropivacaine 0.1% and 0.2% with bupivacaine 0.2% for single-shot caudal anaesthesia in children. *Paediatr Anaesth* 2000, 10:449-504.
22. Taylor MS. Managing of postoperative pain. *Hosp Med* 2001; 62: 560-3.