

1 1 ( )  
1 1

**D**

\*

---

*D*

*D*

*D*

)*D*

*D*

*D*

*D*

( )


*D*

**D**

---

( )

---



1

McNemar's

Ca=2.5-5 mg/dl

$$P = 2.5-5 \text{ mg/dl}$$

(% )

$$l \pm l$$

(% )

Alb=3.5-5.2 gr/dl

D

D

)

(50<sup>s</sup> polydoros

McNemar's

(P < / )

(% / )

mg/kg/day

( )

Roche

/ mg/kg/day

/ mg/kg/day

mg/kg

PTH

D

(% )

(% / )

D

.(P < / )

(P< / )

(% / )

McNemar's

D

D

<b>1,25 VitD</b>	<b>25 VitD</b>	<b>PTH</b>	<b>P</b>	<b>Ca</b>
/ ± /	/ ± /	/ ± /	/ ± /	/ ± /
/ ± /	/ ± /	/ ± /	/ ± /	/ ± /

( ) Leachman

D

D

( )

D

D

D

cAMP

calcium load

D

( ) Margolis

( )

Rodland

D

Margolis

D

D

D

mg/kg

Rodland

(P&lt; / )

( )

Kindmark

D

D

(P&lt; / )


( P&lt; / )

D

Rodland

( )

turnover



## REFERENCES

---

1. Simpson NB, Cunliffe WJ. Disorders of the sebaseous glands In: Burns T, Breathnach S, Cox N, Griffiths C, editors. Rooks text book of dermatology. 7<sup>th</sup> ed. Oxford: Blackwell publishing. 2004; P: 43/1-43/78.
2. Goulden V, Clark SM, McGeown C, Cunliffe WJ. Treatment of acne with intermittent isotretinoin. Br J Dermatol 1997; 137: 106-8.
3. Digiovanna JJ. Isotretinoin effects on bone. J Am Acad Dermatol 2001; 45(5): S 176-82.
4. Pittsley RA, Yoder FW. Retinoid hyperostosis: Skeletal toxicity associated with long-term administration of 13-cis retinoic acid for refractory ichthyosis. N Engl J Med 1983; 306-1012.
5. Ellis CN, Madison KC, Pennes DR, Martel W, Voorhees JJ. Isotretinoin therapy is associated with early skeletal radiographic changes. J Am Acad Dermatol 1984; 10: 1024-9.
6. Lawson JP, McGuire J. the spectrum of skeletal changes associated with long-term administration of 13-cis-retinoic acid. Skeletal Radiol 1987; 16: 91-7.

7. Melnik B, Glück S, Jungblaut RM, Goerz G. Retrospective radiographic study of skeletal changes after long-term etretinate therapy. *Br J Dermatol* 1987; 116: 207-12.
8. Wilson DJ, Kay V, Charig M, Hughes DG, Creasy TS. Skeletal hyperostosis and extraosseous calcification in patients receiving long-term etretinate (Tigason). *Br J Dermatol* 1988; 119: 597-607.
9. DiGiovanna JJ, Helfgott RK, Gerber LH, Peck GL. Extra spinal tendon and ligament calcification associated with long-term therapy with etretinate. *N Engl J Med* 1986; 315: 1177-82.
10. Halkier-Sorensen L, Andresen J. A retrospective study of bone changes in adults treated with etretinate. *J Am Acad Dermatol* 1989; 20: 83-7.
11. White SI, Mackie RM. Bone changes associated with oral retinoid therapy. *Pharmacol Ther* 1989; 40: 137-44.
12. Wilson DJ, Kay V, Charig M, Hughes DG, Creasy TS. Skeletal hyperostosis and extraosseous calcification in patients receiving long-term etretinate(Tigason). *Br J Dermatol* 1988; 119(5): 597-607.
13. Kilcoyne RF, Cope R, Cunningham W, Nardella FA, Denman S, Franz TJ, et al. Minimal spinal hyperostosis with low-dose isotretinoin therapy. *Invest Radiol* 1986; 21: 41-4.
14. Hough S, Avioli LV, Muir H, Gelderblom D, Jenkins G, Kurasi H, et al. Effects of hypervitaminosis A on the bone and mineral metabolism in the rat. *Endocrinology* 1988, 122: 2933-9.
15. Frankel TL, Sechadri MS, McDowall DB, Cornish CJ. Hypervitaminosis A and calcium-regulating hormones in the rat. *J Nutr* 1986; 116: 578-87.
16. Trechsel U, Fleisch H. A new model to study mechanisms of bone resorption in rats using a retinoid. *Calcify Tissue Int* 1984; 36(suppl 2): 549.
17. Rodland O, Aksnes K, Nilsen A, Morken T. Serum level of vit D metabolities in isotretinoin-treated acne patients. *Acta Derm Venred* 1992; 72:217-9.
18. Trechsel U, Fleisch H. Effect of a retinoid on Ca and Vit D metabolism in thyroparathyroidectomized (TPTX) rats. In: Norman AW, Schaefer K, Grigoleit HG, Herrath DV, eds. Vitamin D, chemical, Biochemical and clinical update. Berlin: de gruyter. 1985; P:51.
19. Leachman SA, Insogna KL, Katz L, Ellison A, Milstone LM. Bone densities in patients receiving isotretinoin for cystic acne. *Arch Dermatol* 1999; 135: 961-5.
20. Margolis DJ, Attie M, Leyden JJ. Effects of isotretinoin on bone mineralization during routine therapy with isotretinoin for acne vulgaris. *Arch Dermatol* 1996; 132 (7): 769-74.
22. Kindmark A, Rollmand O, Mallmin H, Petern-Mallmin M, L junghall S, Melhus H. Oral isotretinoin therapy in severe acne induces transient suppression of biochemical markers of bone turnover and calcium homeostasis. *Acta Derm Venereol* 1998; 78 (4): 266-9.