

//
//

()

D

*

D

D

D

)*D*

(*D*

D

D

()

D

D

()

()

()

()

A

(DISH)

()

()

()

()

D

D

()

D

D

D

ELISA

D

•

D

DRG

1,25 (OH)₂ VitD: 39-139 pm

25 OH VitD: 47-144 nm

ELISA

•

DRG

(before and after treatment)

PTH=8.8-76.6 pg/ml

McNemar's

Ca=2.5-5 mg/dl :

P= 2.5-5 mg/dl :

(%)

/ ± /

(%)

Alb=3.5-5.2 gr/dl :

D

D

)

(50^S polydoros

McNemar's

(P < /)

(% /)

mg/kg/day ()

Roche

/ mg/kg/day

(P < /)

mg/kg

PTH

D

(%)

(% /)

D

(P < /)

(P < /)

(% /)

McNemar's

D

D

1,25 VitD	25 VitD	PTH	P	Ca
/ ± /	/ ± /	/ ± /	/ ± /	/ ± /
/ ± /	/ ± /	/ ±	/ ± /	/ ± /

() Leachman

D

D

()

D

D

D

cAMP

D

calcium load

()

Margolis

()

Rodland

D

Margolis

(P< /) D

D

D

D

mg/kg

Rodland

(P< /)

()

Kindmark

D

(P< /)

(P< /)

D

D

Rodland

()

turnover

	/	/			
			t-test	D	
				()	D
					α
Rodland)		()	Trechsew ()
	(()			
					D
				D	
				()	Frankel

REFERENCES

1. Simpson NB, Cunliffe WJ. Disorders of the sebaceous glands In: Burns T, Breathnach S, Cox N, Griffiths C, editors. Rooks text book of dermatology. 7th ed. Oxford: Blackwell publishing. 2004; P: 43/1-43/78.
2. Goulden V, Clark SM, McGeown C, Cunliffe WJ. Treatment of acne with intermittent isotretinoin. Br J Dermatol 1997; 137: 106-8.
3. Digiovanna JJ. Isotertinoin effects on bone. J Am Acad Dermatol 2001; 45(5): S 176-82.
4. Pittsley RA, Yoder FW. Retinoid hyperostosis: Skeletal toxicity associated with long-term administration of 13-cis retinoic acid for refractory ichthyosis. N Engl J Med 1983; 306-1012.
5. Ellis CN, Madison KC, Pennes DR, Martel W, Voorhees JJ. Isotretinoin therapy is associated with early skeletal radiographic changes. J Am Acad Dermatol 1984; 10: 1024-9.
6. Lawson JP, McGuire J. the spectrum of skeletal changes associated with long-term administration of 13-cis-retinoic acid. Skeletal Radiol 1987; 16: 91-7.

7. Melnik B, Glüch S, Jungblaut RM, Goerz G. Retrospective radiographic study of skeletal changes after long-term etretinate therapy. *Br J Dermatol* 1987; 116: 207-12.
8. Wilson DJ, Kay V, Charig M, Hughes DG, Creasy TS. Skeletal hyperostosis and extraosseous calcification in patients receiving long-term etretinate (Tigason). *Br J Dermatol* 1988; 119: 597-607.
9. DiGiovanna JJ, Helfgott RK, Gerber LH, Peck GL. Extra spinal tendon and ligament calcification associated with long-term therapy with etretinate. *N Engl J Med* 1986; 315: 1177-82.
10. Halkier-Sorensen L, Andresen J. A retrospective study of bone changes in adults treated with etretinate. *J Am Acad Dermatol* 1989; 20: 83-7.
11. White SI, Mackie RM. Bone changes associated with oral retinoid therapy. *Pharmacol Ther* 1989; 40: 137-44.
12. Wilson DJ, Kay V, Charig M, Hughes DG, Creasy TS. Skeletal hyperostosis and extraosseous calcification in patients receiving long-term etretinate(Tigason). *Br J Dermatol* 1988; 119(5): 597-607.
13. Kilcoyne RF, Cope R, Cunningham W, Nardella FA, Denman S, Franz TJ, et al. Minimal spinal hyperostosis with low-dose isotretinoin therapy. *Invest Radiol* 1986; 21: 41-4.
14. Hough S, Avioli LV, Muir H, Gelderblom D, Jenkins G, Kurasi H, et al. Effects of hypervitaminosis A on the bone and mineral metabolism in the rat. *Endocrinology* 1988, 122: 2933-9.
15. Frankel TL, Sechadri MS, McDowall DB, Cornish CJ. Hypervitaminosis A and calcium-regulating hormones in the rat. *J Nutr* 1986; 116: 578-87.
16. Trechsel U, Fleisch H. A new model to study mechanisms of bone resorption in rats using a retinoid. *Calcify Tissue Int* 1984; 36(suppl 2): 549.
17. Rodland O, Aksnes K, Nilsen A, Morken T. Serum level of vit D metabolites in isotretinoin-treated acne patients. *Acta Derm Venred* 1992; 72:217-9.
18. Trechsel U, Fleisch H. Effect of a retinoid on Ca and Vit D metabolism in thyroparathyroidectomized (TPTX) rats. In: Norman AW, Schaefer K, Grigoleit HG, Herrath DV, eds. *Vitamin D, chemical, Biochemical and clinical update*. Berlin: de gruyter. 1985; P:51.
19. Leachman SA, Insogna KL, Katz L, Ellison A, Milstone LM. Bone densities in patients receiving isotretinoin for cystic acne. *Arch Dermatol* 1999; 135: 961-5.
20. Margolis DJ, Attie M, Leyden JJ. Effects of isotretinoin on bone mineralization during routine therapy with isotretinoin for acne vulgaris. *Arch Dermatol* 1996; 132 (7): 769-74.

22. Kindmark A, Rollmand O, Mallmin H, Petern-Mallmin M, Ljunghall S, Melhus H. Oral isotretinoin therapy in severe acne induces transient suppression of biochemical markers of bone turnover and calcium homeostasis. *Acta Derm Venereol* 1998; 78 (4): 266-9.