

// ()
 // , ,

*

Visual Analogue Scale

()

Mann-Withney U

) (P < 0/001)

(P< 0/005)

.(

)

(

H₁

.()

()

.()

.()

f-malekzad@yahoo.com

Heyer.()

c

.()

()

()

.()

.()

()

.()

(Revia)

()

(

)

.()

.()

(-)

)

(

.()

.()

Visual Analogue Scale (VAS)

)

VAS

(Visual Analogue Scale) VAS

Margus

.()

(P < /)

(P < /)

Heyer

alloknesis

.()

()

Peer

()

Metze

()

Wotfhegen

REFERENCES

- 1.Chompion RH, Burton JL, Burns DA, Breathnach SM. Textbook of Dermatology. Blackwell Science Ltd. 1998; P:617-27.
- 2.Heyer G, Groene D, Martus P. Efficacy of Naltrexone on Acetylcholine induced alloknesis in atopic eczema. Exp Dermatol 2002; 11: 448-55.

-
3. Metze D, Reimann S, Beissert S, Luger T. Efficacy and safety of Naltrexone , an oral opiate receptor antagonist, in the treatment of pruritus in internal and dermatological diseases. *J Am Acad Dermatol*. 1999; 41: 533-9.
 4. Panli-Margus C, Mikus G, Alscher DM, Kirschner T, Negel W. Naltrexone dose not relieves uremic pruritus: result of a randomised double-blind, placebo-controlled cross over study. *J Am Soc Nephrol* 2000; 11: 514-9.
 5. Peer G, Kivity S, Agami O, Fireman E, Silverberg D, Blum M. Randomized cross over trial of Naltrexone in uremic pruritus. *Lancet* 1996; 348: 1552-54.
 6. Sadock JB, Sadock AV. *Comprehensive Textbook of Psychiatry*. Lippincott. 2000; P:2407-11.
 7. Heyer G, Dotzer M, Diepgen TL, Handwerker H. Opiate and H₁ antagonist effects on histamine induced pruritus and alopecia. *Pain* 1997; 73: 239-43.
 8. Wolfhagen FH, Sternieri E, Hope WC, Vitale G, Bertolotti M, Van Buuren HR. Oral Naltrexone treatment for cholestatic pruritus: a double-blind placebo-controlled study. *Gastroenterology* 1997;113(4):1264-9.