

()

//
//
//

*

Stomacher

BPW

ml

RV ml

/ ml

HE

(% /)

(% /)

(% /) S. thompson

% /

% /

()

)

(

()

(CDC)

()

(food borne diseases)

()

()

()

HACCP

()

% / % /

% / % /

()

% / % /

()


()

()

()

-

(mcg) (mcg)
 (mcg) (mcg) () (ISO 6579)
 (mcg) (mcg) gr
 (mcg) (mcg) ml (Stomacher)
 (mcg) (mcg) (BPW. Merck)
 (mcg) (mcg)
 Stomacher (Seward 400)
 / ml °C
 ml
 Rappaport-Vassiliadis Broth (RV. Merck)
 °C
 (% /)
 (% /)
 () (% /) (HE. Merck)
 °C



Disk Diffusion

()
 (Heart Infusion Broth. Difco)
 °C
 Heart Infusion Broth
 /

(% /) *S. enteritidis* (% /) *S. thompson*
 (% /) *S. paratyphi C*
 % /

(Hi Media)
 (mcg) :

(S. untypable)

.()

.()

()

S.thompson
 S.paratyphi C
 S.meleagridis
 S.enteritidis
 S.virginia
 S.group II
 S.haardt
 S.anatum
 S.veyle
 S.typhimurium
 S.untypable

S (%)	I (%)	R (%)	S (%)	I (%)	R (%)
(/)		(/)	(/)	(/)	(/)
(/)	(/)	(/)	(/)	(/)	()
(/)	(/)	(/)	(/)		(/)
(/)	(/)	()	(/)	(/)	(/)
(/)		(/)	()		
(/)		(/)	()		
()			()		
()			()		
()			()		
()			()		
()			()		
()			()		
()			()		
()			()		
()			()		

S.meleagridis S.paratyphie C S.thompson
 S.haardt S.groupII S.virginia S.enteritidis (/)% /
 S.typhimurium S.veyle S.anatum
 S.thompson
 (P= /)
 pH
 pH
 ()
 S.haardt S.enteritidis
 S.typhimurium S.virchow S.newport
 () S.heidelberg (/)% /
 S.galinarom S.pulorum (% /) (% /)
 () S.cholorasuis S.typhi () (% /)
 S.anatum % /
 S.meleagridis % / ()
 ()
 Luu
 S.haardt S.enteritidis % / Stevens
 S.anatum %
 S.meleagridis % / % /
 S.meleagridis S.anatum ()
 S.enteritidis

() % /

% / % /) ()

% %

% / (

%

()

()

REFERENCES

1. Todd EC. Epidemiology of foodborne disease, a worldwide review. World Health State O. 1997;50:30-50.
2. O'osterom J. Epidemiological studies and proposed preventive measures in the fight against human Salmonellosis. Int J Food Microbiol 1991;21:779-89.
3. Acha. PN and Szyfree B. Zoonoses and communicable diseases common to man and animals. 3rd ed. Washington DC: Pan American Health Organisation. 2001; P:233-46.
4. Health Protection Agency, Centre for Infections. Communicable Disease and health protection quarterly reviews: January to March 2005. J Public Health 2006;27(3):303-7.
5. Bean NH, Griffin PM. Foodborne disease outbreaks in United States 1973-1987: pathogens, vehicles and trends. J Food Prot 1990;53:804-7.
6. White PL, Naugle AL, Jackson CR, Fedorka-Cray PJ, Rose BE. *Salmonella enteritidis* in meat, poultry, and pasteurized egg products regulated by the U.S. Food Safety and Inspection Service, 1998 through 2003; J Food Prot. 2007 Mar;70(3):582-91.

-
7. D' Aoust JY. *Salmonella* and the international food trade. Int J Food Microbiol 1994;24:11-31.
 8. Wray C, Davies RH. *Salmonella* infections in cattle, in *Salmonella* in domestic animals. Oxon, England; Wray C & Wray A, CAB International. 2000; P:169-190.
 9. Bryan FL, Doyle MP. Health risks and consequences of *Salmonella* and *Campylobacter jejuni* in raw poultry. J Food Prot 1995;58:326-44.
 10. Adesiyon AA and Oni OO. Prevalence and antibiograms of *salmonella* in slaughtered cattle, slaughter areas and effluents in Zaria abattoirs, Nigeria; J.Food.Prot 1989; 52:232-235.
 11. Roof MB, Doitchinoff DD. Safety, efficacy, and duration of immunity induced in swine by use of an avirulent live *Salmonella choleraesuis*-containing vaccine. Am J Vet Res 1995;56(1):39-44.
 12. Campo P. Evaluation of susceptibility of *non-typhi Salmonella* in a Spanish hospital (1992-1994) and report *Salmonella* serotype typhimurium isolate resistant to quinolones. Eur J Epidemiol 1997;13:239-41.
 13. Molla B, Alemayehu D, Salah W. Sources and distribution of *Salmonella* serotypes isolated from food animals, slaughterhouse personnel and retail meat products in Ethiopia.1997-2002. Ethiop J Health Dev 2003;17(1):63-70.
 14. Tauxe RV. *Salmonella*: a postmodern pathogen. J Food Prot 1991;54:563-8.
 15. Glynn MY, Bopp C, Dewitt W, Dabney P, Mokhtar M, Angulo FJ. Emergence of multidrug-resistant *Salmonella enterica* serotype typhimurium DT104 infections in the United States. N Engl J Med 1998;338(19):1333-8.
 16. Van Duijkeren E, Houwers DJ. A critical assessment of antimicrobial treatment in uncomplicated *Salmonella enteritidis*. Vet Microbiol 2000;73:61-75.
 17. Angkititrakul S, Chomvarin C, Chaita T, Kanistanon K, Waethewutajarn S. Epidemiology of antimicrobial resistance in *Salmonella* isolated from pork, chicken meat and humans in Thailand. Southeast Asian J Trop Med Public Health 2005;36(6):1510-5.
 18. Wegner HC, Arestrup FM, Gerner- Smidt P. Transfer of antibiotic resistant bacteria from animal to man. Acta Vet Scand 1999;92:51-7.
 19. Molla B, Mesfin A. Alemayehu D. Multiple antimicrobial-resistant *Salmonella* serotypes isolated from chicken carcass and giblets in Debre Zeit and Addis Ababa, Ethiopia. Ethiop J Health Dev 2003;17(2):131-49.
 20. Bohaychuk VM, Gensler GE, King RK, Manninen KI, Sorensen O. Occurrence of pathogens in raw and ready-to-eat meat and poultry products collected from the retail marketplace in Edmonton, Alberta, Canada. J Food Prot 2006;69(9):2176-82.
 21. ISO/ CD 6579 – Microbiology of food and animal feeding stuffs – Horizontal method for the detection of *Salmonella*. 1993. (Revision of ISO 6579: 1993).
 22. www.cdc.gov/lab/disk diff.htm, Disk Diffusion Susceptibility Testing .2007
 23. Jay James M. Modern Food Microbiology, 6th ed. Aspen publisher. 2000.

24. Maharjan M, Joshi V, Joshi DD, Manandhar P. Prevalence of *Salmonella* species in various raw meat samples of a local market in Kathmandu. *Ann N Y Acad Sci* 2006;1081: 249-56.
25. Ha TA, Pham TY. Study of *Salmonella*, *Campylobacter*, and *Escherichia coli* contamination in raw food available in factories, schools, and hospital canteens in Hanoi, Vietnam. *Ann N Y Acad Sci* 2006;1081:262-5.
26. Stevens A, Kaboré Y, Perrier-Gros-Claude JD, Millemann Y, Brisabois A, Catteau M, et al. Prevalence and antibiotic-resistance of *Salmonella* isolated from beef sampled from the slaughterhouse and from retailers in Dakar (Senegal). *Int J Food Microbiol* 2006;110(2):178-86.
27. Luu QH, Fries R, Padungtod P, Tran TH, Kyule MN, Baumann MP, et al. Prevalence of *Salmonella* in retail chicken meat in Hanoi, Vietnam. *Ann N Y Acad Sci* 2006;1081:257-61.
28. Carraminana JJ, Rota C, Augutin I, Herrera A. High prevalence of multiple resistance to antibiotics in *Salmonella serovars* isolated from poultry slaughterhouse in Spain. *Vet Microbiol* 2004;104(1-2):133-9.
29. White PL, Naugle AL, Jackson CR, Fedorka-Cray PJ, Rose BE, Pritchard KM, et al. *Salmonella enteritidis* in meat, poultry, and pasteurized egg products regulated by the U.S. Food Safety and Inspection Service, 1998 through 2003. *J Food Protec* 2007;70(3):582-91.
30. Gay JM, Rice DH, Steiger JH. Prevalence of faecal *Salmonella* shedding by cull dairy cattle marketed in Washington State. *J Food Prot* 1994;7:195-7.