

//
//

()

*

:

()
experimental :

AO

%

mm

%

III

mm

($P < /$)

:

:

adel.ebrahimpour@yahoo.com :

...

...

...

(AO)

Malgaigne

"

mm

(exact reduction)

Lambotte

()

()

()

%

()

experimental

Odd's Ratio

()

	%	%	(%)	(%)
	%	%		/ ± /
			%	%
(%)		III A	% II	% I
mm	(%)		%	III B %
			%	
mm		%	%	%
()				

O.R	(n=) mm	(n=)	
	(%)	(%)	
P< /	(/)	(/)	I
	(/)	(/)	II
	(/)	()	IIIA
	(/)	(/)	IIIB
P< /	(/)	()	
	(/)	()	
P< /	(/)	(/)	
	(/)	(/)	
P< /	(/)	(/)	
		(/)	
		(/)	

% /

()

(P< /) ()

III

(P< /) (Pin tract infection)

(P< /)

()

	(n=)	(n=)
--	-------	-------

P< /	(/)	II I	mm)	(
	(/)	IIIB IIIA		

P< /	(/)	(%)	mm
	(/)		

%

(P< /)

mm

P< /

> mm

(Odd's Ratio=)

III

(B A)

4. Melendez EM, Colon C. Treatment of open tibial fractures with the Orthofix fixator. *Clin Orthop* 1989; 241: 224-30.
5. Lawyer RB, Lubbers LM. Use of the Hoffmann apparatus in the treatment of unstable tibial fractures. *J Bone Joint Surg* 1980; 62A: 1264-73.
6. Schmidt A, Rorabeck CH. Fractures of the tibia treated by flexible external fixation. *Clin Orthop* 1983; 178: 162-72.
7. Egger EL, Gottsauner –Wolf F, Palmar J, Aro HT, Chao EYS. Effects of axial dynamization on bone healing. *J Orthop Trauma* 1993; 34: 185-92.
8. Helland P, Boe A, Molster AO. Open tibial fractures treated with the Ex-fi-re external fixation system. *Clin Orthop* 1996; 326: 209-20.
9. Henley MB, Chapman JR, Agel J, Harvey EJ, Whorton AM, Swiontkowski MF. Treatment of the type II, IIIa and IIIb open fractures of the tibial shaft: a prospective comparison of unreamed interlocking intramedullary nails and half-pin external fixators. *J Orthop Trauma* 1998; 12: 1-7.
10. Kenwright J, Richardson JB, Cunningham JL, White SH, Goodship AE, Adams MA, et al. Axial movement in tibial fractures. A controlled randomised trial of treatment. *J Bone Joint Surg Br* 1995; 77B: 645-8.